

PPH Initial management bundle for Home birth

**BLOOD LOSS
500-1000MLS
&
NO SIGNS OF SHOCK**

NB: All women with blood loss over 500mls require transfer to an obstetric unit for further assessment

DATE: _____

TIME: _____

Identify situation and communicate to colleague: PPH	Time	initial
Request birth partner to call 999 ambulance (use emergency card for birth partner)		
Lie mother flat/warm blankets and reassure		
Commence rubbing up contraction/massaging uterus		
Administer Oxytocin 10iu IM		
Assign scribe if possible: birth partner or colleague		
Consider cannulation		
Administer 2 nd dose Oxytocin 10iu IM or IV if cannulated. *Can still be give if placenta remains in situ and there is active bleeding.		
Consider Tissue: Trauma: Thrombin		
Check placenta and membranes (if delivered.)		
Check for tears: suture if appropriate.		
Misoprostol 1g		
Empty bladder & catheterise with indwelling catheter		
Ergometrine 500mcg IM		
Internal bimanual compression		
Give Hartmann's solution 1ltr IV		
O2 @ 15ltrs high flow		
Record observations		
<ul style="list-style-type: none"> NB: if placenta has not delivered following 1st dose of Oxytocin and CCT do not attempt MROP, transfer to Obstetric unit urgently for medical review. 		
<ul style="list-style-type: none"> SBAR HANDOVER TO AMBULANCE/LABOUR WARD 		