

# AAGBI Safety Guideline

## Management of Severe Local Anaesthetic Toxicity



<b>1</b> <b>Recognition</b>	<b>Signs of severe toxicity:</b> <ul style="list-style-type: none"> <li>• Sudden alteration in mental status, severe agitation or loss of consciousness, with or without tonic-clonic convulsions</li> <li>• Cardiovascular collapse: sinus bradycardia, conduction blocks, asystole and ventricular tachyarrhythmias may all occur</li> <li>• Local anaesthetic (LA) toxicity may occur some time after an initial injection</li> </ul>		
<b>2</b> <b>Immediate management</b>	<ul style="list-style-type: none"> <li>• Stop injecting the LA</li> <li>• Call for help</li> <li>• Maintain the airway and, if necessary, secure it with a tracheal tube</li> <li>• Give 100% oxygen and ensure adequate lung ventilation (hyperventilation may help by increasing plasma pH in the presence of metabolic acidosis)</li> <li>• Confirm or establish intravenous access</li> <li>• Control seizures: give a benzodiazepine, thiopental or propofol in small incremental doses</li> <li>• Assess cardiovascular status throughout</li> <li>• Consider drawing blood for analysis, but do not delay definitive treatment to do this</li> </ul>		
<b>3</b> <b>Treatment</b>	<table border="1"> <tr> <td data-bbox="513 1030 1034 1712"> <b>IN CIRCULATORY ARREST</b> <ul style="list-style-type: none"> <li>• Start cardiopulmonary resuscitation (CPR) using standard protocols</li> <li>• Manage arrhythmias using the same protocols, recognising that arrhythmias may be very refractory to treatment</li> <li>• Consider the use of cardiopulmonary bypass if available</li> </ul> <b>GIVE INTRAVENOUS LIPID EMULSION</b>                      (following the regimen overleaf)                     <ul style="list-style-type: none"> <li>• Continue CPR throughout treatment with lipid emulsion</li> <li>• Recovery from LA-induced cardiac arrest may take &gt;1 h</li> <li>• Propofol is not a suitable substitute for lipid emulsion</li> <li>• Lidocaine should not be used as an anti-arrhythmic therapy</li> </ul> </td><td data-bbox="1034 1030 1550 1712"> <b>WITHOUT CIRCULATORY ARREST</b>                      Use conventional therapies to treat:                     <ul style="list-style-type: none"> <li>• hypotension,</li> <li>• bradycardia,</li> <li>• tachyarrhythmia</li> </ul> <b>CONSIDER INTRAVENOUS LIPID EMULSION</b>                      (following the regimen overleaf)                     <ul style="list-style-type: none"> <li>• Propofol is not a suitable substitute for lipid emulsion</li> <li>• Lidocaine should not be used as an anti-arrhythmic therapy</li> </ul> </td></tr> </table>	<b>IN CIRCULATORY ARREST</b> <ul style="list-style-type: none"> <li>• Start cardiopulmonary resuscitation (CPR) using standard protocols</li> <li>• Manage arrhythmias using the same protocols, recognising that arrhythmias may be very refractory to treatment</li> <li>• Consider the use of cardiopulmonary bypass if available</li> </ul> <b>GIVE INTRAVENOUS LIPID EMULSION</b> (following the regimen overleaf) <ul style="list-style-type: none"> <li>• Continue CPR throughout treatment with lipid emulsion</li> <li>• Recovery from LA-induced cardiac arrest may take &gt;1 h</li> <li>• Propofol is not a suitable substitute for lipid emulsion</li> <li>• Lidocaine should not be used as an anti-arrhythmic therapy</li> </ul>	<b>WITHOUT CIRCULATORY ARREST</b> Use conventional therapies to treat: <ul style="list-style-type: none"> <li>• hypotension,</li> <li>• bradycardia,</li> <li>• tachyarrhythmia</li> </ul> <b>CONSIDER INTRAVENOUS LIPID EMULSION</b> (following the regimen overleaf) <ul style="list-style-type: none"> <li>• Propofol is not a suitable substitute for lipid emulsion</li> <li>• Lidocaine should not be used as an anti-arrhythmic therapy</li> </ul>
<b>IN CIRCULATORY ARREST</b> <ul style="list-style-type: none"> <li>• Start cardiopulmonary resuscitation (CPR) using standard protocols</li> <li>• Manage arrhythmias using the same protocols, recognising that arrhythmias may be very refractory to treatment</li> <li>• Consider the use of cardiopulmonary bypass if available</li> </ul> <b>GIVE INTRAVENOUS LIPID EMULSION</b> (following the regimen overleaf) <ul style="list-style-type: none"> <li>• Continue CPR throughout treatment with lipid emulsion</li> <li>• Recovery from LA-induced cardiac arrest may take &gt;1 h</li> <li>• Propofol is not a suitable substitute for lipid emulsion</li> <li>• Lidocaine should not be used as an anti-arrhythmic therapy</li> </ul>	<b>WITHOUT CIRCULATORY ARREST</b> Use conventional therapies to treat: <ul style="list-style-type: none"> <li>• hypotension,</li> <li>• bradycardia,</li> <li>• tachyarrhythmia</li> </ul> <b>CONSIDER INTRAVENOUS LIPID EMULSION</b> (following the regimen overleaf) <ul style="list-style-type: none"> <li>• Propofol is not a suitable substitute for lipid emulsion</li> <li>• Lidocaine should not be used as an anti-arrhythmic therapy</li> </ul>		
<b>4</b> <b>Follow-up</b>	<ul style="list-style-type: none"> <li>• Arrange safe transfer to a clinical area with appropriate equipment and suitable staff until sustained recovery is achieved</li> <li>• Exclude pancreatitis by regular clinical review, including daily amylase or lipase assays for two days</li> <li>• Report cases as follows:                         <ul style="list-style-type: none"> <li>in the United Kingdom to the National Patient Safety Agency (via <a href="http://www.npsa.nhs.uk">www.npsa.nhs.uk</a>)</li> <li>in the Republic of Ireland to the Irish Medicines Board (via <a href="http://www.imb.ie">www.imb.ie</a>)</li> </ul> </li> </ul> <p>If Lipid has been given, please also report its use to the international registry at <a href="http://www.lipidregistry.org">www.lipidregistry.org</a>. Details may also be posted at <a href="http://www.lipidrescue.org">www.lipidrescue.org</a></p>		

**Your nearest bag of Lipid Emulsion is kept.....**

This guideline is not a standard of medical care. The ultimate judgement with regard to a particular clinical procedure or treatment plan must be made by the clinician in the light of the clinical data presented and the diagnostic and treatment options available.

© The Association of Anaesthetists of Great Britain & Ireland 2010

## IMMEDIATELY

Give an initial intravenous bolus injection of 20% lipid emulsion  
 $1.5 \text{ ml.kg}^{-1}$  over 1 min

AND

Start an intravenous infusion of 20% lipid emulsion at  $15 \text{ ml.kg}^{-1}.\text{h}^{-1}$

## AFTER 5 MIN

Give a **maximum of two** repeat boluses (same dose) if:

- cardiovascular stability has not been restored or
- an adequate circulation deteriorates

Leave **5 min** between boluses

A maximum of **three** boluses can be given (including the initial bolus)

AND

Continue infusion at same rate, but: **Double** the rate to  $30 \text{ ml.kg}^{-1}.\text{h}^{-1}$  at any time after 5 min, if:

- cardiovascular stability has not been restored or
- an adequate circulation deteriorates

Continue infusion until stable and adequate circulation restored or maximum dose of lipid emulsion given

***Do not exceed a maximum cumulative dose of  $12 \text{ ml.kg}^{-1}$***

***An approximate dose regimen for a 70-kg patient would be as follows:***

## IMMEDIATELY

Give an initial intravenous bolus injection of 20% lipid emulsion  
100 ml over 1 min

AND

Start an intravenous infusion of 20% lipid emulsion at  $1000 \text{ ml.h}^{-1}$

## AFTER 5 MIN

Give a **maximum of two** repeat boluses of 100 ml

AND

Continue infusion at same rate but **double** rate to  $2000 \text{ ml.h}^{-1}$  if indicated at any time

***Do not exceed a maximum cumulative dose of 840 ml***



This AAGBI Safety Guideline was produced by a Working Party that comprised:  
Grant Cave, Will Harrop-Griffiths (Chair), Martyn Harvey, Tim Meek, John Picard, Tim Short and Guy Weinberg.

**This Safety Guideline is endorsed by the Australian and New Zealand College of Anaesthetists (ANZCA).**