

TARGET	Adult Critical Care Unit (ACCU)	
AUDIENCE	University Hospital Wishaw	
PATIENT GROUP	Patients in ACCU	

Clinical Guidelines Summary

Metaraminol is a vasoconstrictor sympathomimetic, raising blood pressure by acting on alpha-adrenoceptors to constrict peripheral vessels. It can be used as continuous infusion for the treatment of mild/moderate hypotension in association with an Epidural in ACCU in the postoperative period.

This guideline informs in 3 sections about

- 1. Reconstitution and administration
- 2. Prescribing, dosing and monitoring
- 3. Adverse effects



Reconstitution and Administration

Standard concentration is 0.5mg/ml. Dilute two 10mg vials Metaraminol with 38ml sodium chloride 0.9% Normal Saline in a 50ml Syringe. Use the largest vein present unless a central line is available. The cannula should be checked carefully for extravasation prior to starting the infusion. If there is any doubt about the cannula medical review should be sought.

Prescribing, dosing and monitoring

This should be prescribed on HEPMA as Metaraminol-infusion as charted. Non-invasive pressure should be checked every 5 minutes until stable thereafter every hour. Standard Epidural Checks/Obs should be carried out as normal. An Arterial Line is not necessary. The cannula should be checked HOURLY to ensure blanching/extravasation does not occur as this can result in skin necrosis. The infusion rate can be changed and titrated to a MAP of >65mm Hg or higher if specified by the anaesthetist. Range of rate is 0 – 20ml/hr. Any higher requirements should prompt immediate review by the prescribing anaesthetist or (if out of hours or not available) by the ACCU-Trainee. The patient must be fully assessed and other causes of hypotension (bleeding, sepsis) actively sought. Avoid abrupt withdrawal. The infusion should not be stopped suddenly but gradually titrated off. In general the duration of the Metaraminol-Infusion should not exceed the duration of the Epidural.

Lead Author	Stephan Dalchow, Jennifer Murphy	Date approved	6.12.24
Version	3	Review Date	1.1.30



Adverse effects

If a patient becomes hypertensive stop the infusion immediately. If the blood pressure is dangerously high and not coming down quickly – contact an anaesthetist urgently. Hypertension can result in reflex bradycardia. If this occurs and does not resolve following discontinuation of the infusion it can be treated by a bolus of 200 micrograms of glycopyrrolate or 600 micrograms of atropine.

Lead Author	Stephan Dalchow, Jennifer Murphy	Date approved	6.12.24
Version	3	Review Date	1.1.30



Lead Author	Stephan Dalchow, Jennifer Murphy	Date approved	6.12.24
Version	3	Review Date	1.1.30



Appendices

1. Governance information for Guidance document

Lead Author(s):	Stephan Dalchow; Jennifer Murphy
Endorsing Body:	ADTC
Version Number:	3
Approval date	6.12.24
Review Date:	1.1.2030
Responsible Person (if different from lead author)	

CONSULTATION AND DISTRIBUTION RECORD		
Contributing Author / Authors Renewal Stephan Dalchow, Jennifer Murphy		
Consultation Process / Stakeholders:		

Lead Author	Stephan Dalchow, Jennifer Murphy	Date approved	6.12.24
Version	3	Review Date	1.1.30



Distribution	

CHANGE RECORD				
Date	Lead Author	Change Version No.		
		e.g. Review, revise and update of policy in line with contemporary professional structures and practice	1	
			2	
6.12.2024	Stephan Dalchow	Put in requested format Wishaw General Hospital replaced by University Hospital Wishaw Medicines chart replaced by HEPMA	3	
			4	
			5	

Lead Author	Stephan Dalchow, Jennifer Murphy	Date approved	6.12.24
Version	3	Review Date	1.1.30



Lead Author	Stephan Dalchow, Jennifer Murphy	Date approved	6.12.24
Version	3	Review Date	1.1.30