Seizure/Fits

Observations/NEWS

Read Patient Alerts



Temperature

Pulse

Respiratory rate

Oxygen saturations (on air or O2 %)

Blood glucose

✓ AVPU✓ Pupil size

Market Consider Alcometer

Complete CIWA/GMAWS if ?alcohol Withdrawal seizure

Investigations

<u>Discuss with senior staff; not all patients will require all investigations, especially patients with known seizure disorders</u>

If NO history of Epilepsy/or alcohol excess

Consider:

- · IV access
- Admission bloods (as per order set): FBC, U&E, LFTs, add Glucose, Ca, Mg²⁺, Phosphate & CK.
- · VBG (lactate)
- ECG
- · CT head after discussion with senior

If Pregnant: MEWS chart, BP, urinalysis

Treatments

Consider:

- If blood glucose <4mmol/l follow 'Management of Hypoglycaemia' guidelines (Overleaf)
- · If history of alcohol excess/dependence follow CIWA/GMAWS guidance
- Benzodiazepines (as per CIWA/GMAWS guidelines).
- · IV fluid if BP low

Red flags

- Head injury: Important to establish collateral history, seizure activity could be post traumatic.
- Abnormal ECG: Cardiac arrhythmias and/or electrolyte imbalances can lead to seizure activity.
- History of overdose or poisoning: Ensure Toxbase is checked.
- Pregnancy: Escalate to senior staff MEWS chart should be used.
- Purpura: A rash on any part of the body that is caused by small haemorrhages under the skin. A purpuric rash does not blanch (go white) when pressure is applied to it.
- Non-blanching rash: A rash that does not go return to natural skin colour after pressure is applied.
- Signs of meningism: Classically a stiff neck together with headache and photophobia.

Patient Presentation Cards			Version 1	Page 1 of 1		
Document: PPC's			Author: Dr G Pickerin Hughes	ng, H Godfrey, C	NHS	WEDIC ON