

Appendix 4 - Chlordiazepoxide Fixed Reducing Dose Chart: (Lower Risk)

Addressograph Label

Name:	DoB:	Ward:	CHI:	Consultant:
--------------	-------------	--------------	-------------	--------------------

Day	Date	Drug	Prescribed Dose	Route	Administration Times Given by (initials)				Prescriber's Signature and Name
					0800 hours	1200 hours	1800 hours	2200 hours	
1		Chlordiazepoxide	30mg	Oral	-----	-----	-----	-----	
2		Chlordiazepoxide	20mg	Oral	-----	-----	-----	-----	
3		Chlordiazepoxide	20mg	Oral	-----			-----	
		Chlordiazepoxide	10mg	Oral		-----	-----		
4		Chlordiazepoxide	10mg	Oral	-----		-----	-----	
5		Chlordiazepoxide	10mg	Oral	-----			-----	

PRESCRIBE

And “

On drug chart: “Chlordiazepoxide according to Reducing Dose Chart”.

Chlordiazepoxide 20-50mg as required, as per GMAWS score (250mg maximum, including regular + PRN dose in 24 hours)”; review use daily

CONSIDER

Is this the right chart?

Are nursing staff monitoring with GMAWS?

Has IV vitamin B&C or IM thiamine been started?

Is this regimen being reviewed?

– This chart is for patients in alcohol withdrawal assessed as being at high risk of complications from Benzo's

– Breakthrough symptoms should be assessed using the GMAWS and treated

– All patients in alcohol withdrawal are at risk of Wernicke's Encephalopathy

– Must be reviewed daily and considered if needs adjusted